TEST

Histoplasma complement fixation

A blood test that looks for antibodies made in response to Histoplasma fungus

WHAT IT MEASURES
Antibodies to Histoplasma yeast and mold-phase antigens

SAMPLE
Blood

FASTING NEEDED?
No

HOW LONG IT TAKES
Usually less than five minutes

RESULTS IN
Several days, depending on laboratory and referral testing

USED FOR
Supporting diagnosis of current or recent histoplasmosis

MAIN LIMITATION
Early infection and weak immune responses can cause false-negative results

Overview

Histoplasma complement fixation is a blood test that looks for antibodies to Histoplasma , a fungus found in soil enriched with bird or bat droppings. Antibodies are proteins made by your immune system after it encounters a substance it recognizes as foreign.

The test usually measures antibodies against two forms of the fungus: yeast-phase and mycelial, or mold-phase, antigens. An antigen is a substance that can trigger an immune response.

"Complement fixation" describes the laboratory method. Your serum, the liquid part of blood, is mixed with Histoplasma antigens and laboratory complement proteins. The reaction helps show whether Histoplasma antibodies are present and estimates their amount.

This test does not identify the fungus itself. It is one part of an assessment that may also include urine or blood antigen testing, imaging, cultures, tissue examination, and your history of possible exposure.

Why the test is ordered

The test may help investigate symptoms and imaging findings that could be related to histoplasmosis. These can overlap with many other infections and lung conditions, so an antibody result cannot establish the cause by itself.

A healthcare professional may order complement fixation testing when:

Complement fixation is often ordered with Histoplasma immunodiffusion, another antibody method. Antigen testing may also be ordered because it detects parts of the fungus rather than your immune response.

- You have a compatible illness after possible exposure to contaminated soil
- A chest X-ray or CT scan shows changes that need explanation
- Histoplasmosis is one of several possible causes of a lung infection
- An initial Histoplasma test is negative but suspicion remains
- Comparing two blood samples may help show whether antibody levels are rising

Who should have the test

This test is mainly used for people whose symptoms, exposure history, examination, or imaging suggest a possible Histoplasma infection. Exposure may occur while disturbing contaminated soil, cleaning areas with bird or bat droppings, exploring caves, or working around old buildings.

It is not generally used as a stand-alone screening test for people without symptoms. A positive result can reflect an earlier infection rather than a current illness, especially in places where Histoplasma exposure is common.

People with weakened immune systems may need other tests even if complement fixation is negative. This includes some people taking immune-suppressing medicines or living with conditions that reduce the body's ability to make antibodies.

How to prepare

You do not usually need to fast. You can normally eat, drink water, and take your usual medicines.

Tell the person ordering the test about medicines that suppress the immune system, such as chemotherapy or medicines used after an organ transplant. These medicines may reduce

antibody production and affect how a negative result is interpreted.

Also mention a recent blood-product transfusion or antibody treatment. If other blood tests are being collected at the same appointment, their preparation instructions may be different.

What happens during the test

A healthcare worker places a band around your upper arm and cleans the skin. A small needle is inserted into a vein, usually inside the elbow, and blood is collected into a tube.

The needle is then removed and pressure is placed over the site. The collection itself usually takes less than five minutes, although the full appointment may take longer.

The sample may be sent to a reference laboratory because not every local laboratory performs fungal antibody testing.

Understanding your results

Complement fixation results are usually reported as a titer . A titer shows how far the laboratory could dilute your serum while still detecting antibodies.

For example, a result with a larger second number represents antibody detection at a greater dilution. However, the number should not be interpreted without the laboratory's reference information and your clinical context.

A negative result means the laboratory did not detect antibodies above its reporting threshold. It does not completely rule out histoplasmosis. Antibodies may not yet have developed, or your immune system may not produce a strong antibody response.

A positive result means Histoplasma-reactive antibodies were detected. It may support a recent or current infection, but it can also reflect an older exposure or a reaction to another fungus.

A rising titer can be more informative than one result. When appropriate, a second sample collected later may show a fourfold change-for example, from 1:8 to 1:32. The timing and importance of such a change must be interpreted with your symptoms and other test results.

Ranges and interpretation vary by laboratory, assay method, age, sex, pregnancy status, medicines, immune function, and clinical context. Use the reference information printed on your own report.

What affects the results

Several factors can make the result misleading even when the laboratory has performed the test correctly.

Testing too early is a major cause of a negative result. Antibodies often take several weeks to become detectable after infection. Repeat antibody testing or a different type of test may be considered if symptoms and exposure strongly suggest infection.

A weakened immune response can reduce antibody production. Immune-suppressing medicines, organ transplantation, advanced HIV infection, and some blood or immune disorders may make the test less likely to detect an infection.

Previous exposure can leave antibodies in the blood after the illness has resolved. A positive test therefore does not always mean the fungus is currently causing symptoms.

Other fungal infections can produce cross-reactions. Antibodies related to fungi such as Blastomyces may react in Histoplasma tests and cause a positive result when histoplasmosis is not the explanation.

Laboratory methods differ. Laboratories may use different antigens, thresholds, and reporting systems. Results from different laboratories may not be directly comparable.

A recent transfusion or treatment made from donated antibodies can occasionally complicate antibody-test interpretation. Ordinary meals, coffee, exercise, and the time of day are not known to meaningfully change the result.

Accuracy and limitations

Complement fixation cannot reliably confirm or exclude histoplasmosis on its own. Its accuracy changes according to the form and stage of illness and how well your immune system makes antibodies. Because of these differences, one accuracy percentage would not apply to everyone.

The test is more likely to miss an infection when blood is collected early or when immune function is reduced. In these settings, Histoplasma antigen testing, culture, microscopy, or tissue testing may be more useful.

The test may say antibodies are present when histoplasmosis is not the cause of the current illness. This can happen because antibodies remain after an earlier exposure or because another fungal infection produces a cross-reaction.

Antigen testing is often especially useful in widespread, or disseminated, histoplasmosis and in people with weakened immune systems. Culture can directly demonstrate the organism, but

growth may take weeks. Examining tissue or respiratory samples can sometimes provide faster evidence, though obtaining those samples may involve a separate procedure.

A result is most useful when combined with symptoms, exposure history, imaging, and other laboratory findings.

Risks and complications

The risks are those of a routine blood draw. You may have brief discomfort, a small bruise, or feel light-headed; serious complications are uncommon.

What happens next

A negative result may need no further action if your symptoms are improving and other findings are reassuring. If testing was done early, symptoms continue, or immune function is reduced, the next step may be repeat antibody testing or a different test.

A positive result is reviewed alongside your history and other findings. Possible next tests include urine or blood antigen testing, chest imaging, fungal culture, or examination of respiratory fluid or tissue.

Arrange routine follow-up to discuss results that are positive, unclear, or inconsistent with how you feel. Contact a healthcare professional promptly if symptoms are worsening rather than waiting for a scheduled results appointment.

Seek urgent medical care for significant trouble breathing, coughing up more than small streaks of blood, fainting, or rapidly worsening weakness. Call emergency services for severe breathing difficulty, blue or gray lips, new confusion, or severe chest pain.

Common questions

Can I drink coffee before the test?

Yes. Coffee does not normally affect Histoplasma complement fixation. Follow separate instructions if other tests are being collected at the same time.

Does the time of day matter?

No. Antibody titers are not known to change enough during the day to require morning or evening collection.

Can I exercise beforehand?

Yes. Ordinary exercise is not expected to affect the result. Rest before the blood draw if exercise makes you feel light-headed or dehydrated.

How soon after exposure will the test turn positive?

Antibodies can take several weeks to appear. A test collected early may be negative even when further investigation is appropriate.

Does a positive result mean I have active histoplasmosis?

Not necessarily. It may represent current infection, recent infection, older exposure, or a cross-reaction with another fungus. Other findings are needed to determine its meaning.

Why might I also need a urine antigen test?

Antibody and antigen tests look for different things. Antibody testing measures your immune response, while antigen testing looks for parts of the fungus. Using both can improve the assessment in some clinical situations.

Sources

- Centers for Disease Control and Prevention. Clinical Overview of Histoplasmosis .
- Centers for Disease Control and Prevention. Testing Algorithm for Histoplasmosis .
- Wheat LJ, Freifeld AG, Kleiman MB, et al. Clinical practice guidelines for the management of patients with histoplasmosis: 2007 update by the Infectious Diseases Society of America. Clinical Infectious Diseases . 2007;45(7):807-825.
- Azar MM, Hage CA. Laboratory diagnostics for histoplasmosis. Journal of Clinical Microbiology . 2017;55(6):1612-1620.
- Mayo Clinic Laboratories. Histoplasma Antibody Complement Fixation and Immunodiffusion, Serum: Test Catalog and Interpretation .
- ARUP Laboratories. Histoplasma Antibodies by Complement Fixation: Test Directory .