TEST

Quantitative nephelometry test

Quantitative nephelometry measures specific proteins by detecting light scattered from a blood sample

WHAT IT MEASURES
Amount of a specific protein using scattered light

SAMPLE
Usually blood

FASTING NEEDED?
No, unless required for other ordered tests

HOW LONG IT TAKES
About five to ten minutes

RESULTS IN
Usually one to three days, varying by laboratory

USED FOR
Diagnosis and monitoring

MAIN LIMITATION
Very high protein levels can appear falsely low

Overview

Quantitative nephelometry is a laboratory method used to measure the amount of a particular protein in a sample, usually blood. It is not one single test. The name describes how the laboratory makes the measurement.

During testing, the laboratory adds antibodies designed to attach to the protein being measured. The resulting clusters scatter a beam of light. An instrument measures that scattered light and calculates the protein concentration.

Nephelometry may be used to measure:

Your request form or result should name the specific protein. That name matters more for interpretation than the word "nephelometry."

- Immunoglobulins, also called antibodies, such as IgG, IgA, and IgM
- Complement proteins, including C3 and C4
- Certain inflammation-related proteins

- Proteins such as alpha-1 antitrypsin or transferrin
- Free light chains, which are parts of antibodies

Why the test is ordered

The reason for ordering quantitative nephelometry depends on the protein being measured.

Immunoglobulin measurements can help investigate repeated or unusual infections, suspected immune system problems, or unexpectedly high blood protein levels. They may also be used to monitor an established antibody disorder or treatment with immunoglobulin products.

Complement measurements can support the assessment or monitoring of certain inflammatory and autoimmune conditions. Complement proteins form part of the immune system and help antibodies clear microbes and damaged cells.

Other proteins have different purposes. For example, alpha-1 antitrypsin testing may be ordered when investigating certain liver or lung problems. A C-reactive protein measurement can help assess inflammation, although it cannot identify the cause by itself.

Nephelometry often provides one part of an assessment. Your symptoms, medical history, examination, and other test results remain important.

Who should have the test

You may be offered this test if there is a specific reason to measure a blood protein. Examples include repeated infections, possible immune system dysfunction, unexplained inflammation, or monitoring of a known condition.

It may also be ordered after another blood test finds an abnormal total protein level. If an unusual antibody protein is suspected, nephelometry may be combined with serum protein electrophoresis and immunofixation. These tests examine the pattern and type of proteins rather than only their total amount.

Quantitative nephelometry is not a general screening test for everyone. Whether it is useful depends on which protein is being measured and the clinical question.

How to prepare

Fasting is not usually needed. You can normally eat and drink as usual unless the same blood

draw includes another test that requires fasting.

Drink enough water beforehand unless you have been told to restrict fluids. Being well hydrated can make the blood draw easier and reduces the chance that dehydration will concentrate blood proteins.

Tell the healthcare professional arranging the test about:

Do not stop medicines or delay treatment solely because of this test unless you have been given specific instructions.

- Prescription and non-prescription medicines
- Immunoglobulin replacement or antibody-based treatment
- Recent blood or plasma transfusions
- Recent infections, vaccinations, or major inflammation
- Pregnancy or possible pregnancy

What happens during the test

Quantitative nephelometry usually requires a blood sample from a vein in your arm.

A band is placed around your upper arm, and the skin is cleaned. A small needle is inserted into a vein, and blood is collected into a tube. The needle is then removed and pressure is applied to the site.

The blood draw usually takes five to ten minutes. The sample is sent to a laboratory, where the requested protein is measured with a nephelometer.

Less commonly, nephelometry may be used on another body fluid. Collection then depends on the type of sample and will be explained separately.

Understanding your results

Your report should identify the protein measured, its concentration, the units, and the laboratory's reference range. A reference range describes results commonly found in a selected comparison population.

There is no single reference range for "quantitative nephelometry." Ranges differ according to the protein and laboratory method. Interpretation can also vary with age, sex, pregnancy status, medicines, and clinical context.

A high result means more of the measured protein was detected. A low result means less was

detected. The meaning is protein-specific. For example, low immunoglobulin levels and low complement levels have different possible explanations and require different follow-up.

One result rarely establishes a diagnosis. A result just outside the stated range may not represent illness, while a result inside the range may not answer every clinical question. Trends can be useful, particularly when repeat samples are tested by the same laboratory and method.

What affects the results

Several factors can change the true protein level or interfere with the measurement.

Dehydration can make some protein concentrations appear higher because there is less water in the blood. Receiving large amounts of fluid can have the opposite effect.

A recent infection, vaccination, inflammatory episode, surgery, or injury can alter immune and inflammation-related proteins. Strenuous exercise may temporarily affect some inflammatory proteins, but it is not an important influence on every nephelometric test.

Pregnancy and age can change expected protein levels. Kidney or liver function may also matter because these organs help produce, process, or remove many proteins.

Medicines can change the actual level being measured. Steroids like prednisone or hydrocortisone , immune-suppressing medicines, antibody treatments, and immunoglobulin replacement are especially relevant for some tests. A recent transfusion or plasma-containing blood product may also affect antibody and protein measurements.

The sample itself can interfere with scattered-light measurements. Large amounts of fat in the blood can make it cloudy. Breakdown of red blood cells during collection, called haemolysis, may also interfere with some assays. Laboratories commonly check for these problems and may request another sample.

Extremely high protein concentrations can sometimes cause an "antigen excess" or "hook" effect. Too much of the target protein prevents the expected particle clusters from forming, producing a result that is falsely low. Modern instruments use checks and sample dilution to reduce this risk, but it cannot always be excluded.

Accuracy and limitations

Nephelometry can measure proteins precisely when the result falls within the assay's working range and the sample is suitable. However, accuracy differs by protein, instrument, antibody reagent, and laboratory procedure.

There is no single figure for how often quantitative nephelometry misses a problem or gives an abnormal result when no disorder is present. It is a method used for many different proteins, and each assay performs differently.

False-low results may occur with antigen excess, sample dilution errors, or deterioration of the target protein. False-high results can result from cloudy samples or from antibodies reacting with substances similar to the intended target. Laboratories may repeat, dilute, or test a sample by another method when the number does not fit the clinical picture.

Nephelometry tells you how much of a protein is present, but not always what type it is or why it changed. Serum protein electrophoresis and immunofixation are often better for identifying an unusual single antibody protein. Genetic testing or protein phenotyping may be needed for some inherited protein disorders.

Risks and complications

A blood draw may cause brief discomfort, a small bruise, or minor bleeding at the needle site. Some people feel faint, but serious complications are uncommon.

What happens next

A result within the laboratory range may need no further testing, depending on why it was ordered. If symptoms continue, other tests may still be appropriate because a normal protein level does not rule out every possible cause.

An unexpected result is commonly reviewed at a routine follow-up appointment. Possible next steps include repeating the measurement, checking related proteins, reviewing medicines and recent treatments, or arranging electrophoresis, immunofixation, urine testing, or another targeted test.

A nephelometry result by itself rarely indicates an emergency. If the laboratory identifies a result needing prompt review, the requesting service may contact you. Seek urgent medical care based on serious symptoms-such as severe breathing difficulty, fainting, new confusion, or swelling of the tongue or throat-rather than waiting for follow-up of the laboratory number.

Common questions

Can I drink coffee before the test?

Usually, yes. Fasting is not normally required for nephelometry, but coffee may not be allowed if other tests ordered with it require fasting. Follow the instructions for the whole set of tests.

Does the time of day matter?

Usually not. Some proteins can change with inflammation, treatment timing, or other biological factors, so repeat monitoring is sometimes performed under similar conditions.

Can I exercise beforehand?

Ordinary activity is generally acceptable. Strenuous exercise can temporarily alter some inflammation-related proteins, so mention recent intense exercise if that protein is being measured.

How quickly will I get the result?

Many laboratories report routine results within one to three days. Specialist tests or samples sent to another laboratory may take longer.

Is nephelometry the same as protein electrophoresis?

No. Nephelometry measures the amount of a selected protein. Electrophoresis separates proteins into patterns and can help identify an unusual protein.

Why might the test be repeated?

A repeat sample may confirm an unexpected result, check for change over time, or resolve possible interference. The laboratory may also dilute a sample if the protein level is outside the instrument's measuring range.

Sources

- MedlinePlus: Immunoglobulins Blood Test , US National Library of Medicine
- ARUP Consult: Immunoglobulin Disorders , ARUP Laboratories
- Tate J, Ward G. Interferences in Immunoassay . Clinical Biochemist Reviews . 2004;25(2):105-120
- Clinical and Laboratory Standards Institute. Interference Testing in Clinical Chemistry . CLSI guideline EP07, third edition, 2018
- McPherson RA, Pincus MR, editors. Henry's Clinical Diagnosis and Management by Laboratory Methods . 24th edition. Elsevier, 2021
- Rifai N, Horvath AR, Wittwer CT, editors. Tietz Textbook of Laboratory Medicine . Seventh edition. Elsevier, 2023