TEST

TBG blood test

A TBG blood test measures the main protein that carries thyroid hormones through your bloodstream.

WHAT IT MEASURES
Blood level of thyroxine-binding globulin, a thyroid hormone carrier protein

SAMPLE
Blood

FASTING NEEDED?
No

HOW LONG IT TAKES
About 5 to 10 minutes

RESULTS IN
Usually 1 to 3 days, varying by laboratory

USED FOR
Investigating unexpected thyroid hormone test patterns

MAIN LIMITATION
Does not show whether your thyroid gland works normally

Overview

A TBG blood test measures thyroxine-binding globulin. TBG is a protein made mainly by your liver. It carries most of the thyroid hormones called thyroxine, or T4, and triiodothyronine, or T3, through your blood.

Only a small amount of T4 and T3 circulates without a carrier protein. This is called "free" hormone. Free thyroid hormones are available for your body to use.

Changes in TBG can alter total T4 and T3 results without changing the amount of free hormone. For example, a higher TBG level can cause total T4 to rise even when your thyroid gland is working normally.

The TBG test is therefore not a stand-alone test of thyroid function. It is usually used to explain a thyroid test pattern that does not fit your symptoms, medical history, or other results.

Why the test is ordered

Your healthcare professional may order TBG when total T4 or T3 is unexpectedly high or low but thyroid-stimulating hormone, called TSH, and free T4 do not suggest a thyroid disorder.

The test can help investigate:

Testing TBG may prevent a change in carrier protein from being mistaken for an overactive or underactive thyroid.

- An unexpectedly high or low total T4 result
- A possible inherited TBG deficiency or excess
- Changes caused by pregnancy, medicines, or liver disease
- A low total T4 result found during newborn screening
- Differences between your symptoms and thyroid blood test results

Who should have the test

Most people having thyroid tests do not need a TBG measurement. TSH and free T4 are usually more useful for checking how the thyroid gland is working.

TBG testing may be appropriate if your total thyroid hormone results remain unusual or conflict with other tests. It may also be used when a close relative has an inherited TBG difference.

Some newborn screening programs measure total T4. If a baby has a low total T4 but a TSH result that does not suggest an underactive thyroid, TBG testing may help explain the finding. Follow-up testing is still important because TBG deficiency and thyroid disease can occur separately.

How to prepare

You usually do not need to fast. You can generally eat, drink water, and have coffee as normal unless you are having other blood tests that require fasting.

Tell the person who ordered the test if you are pregnant or taking:

These may affect TBG or other thyroid test results. Do not stop prescribed medicines or supplements unless you have been specifically told to do so.

If TBG is being tested with other blood tests, follow the preparation instructions for the full set of tests.

- A combined contraceptive pill
- Oral estrogen or hormone replacement therapy
- Testosterone or anabolic steroids
- Glucocorticoid medicines, such as prednisone
- Thyroid medicine
- High-dose biotin supplements

What happens during the test

A healthcare professional places a small needle into a vein, usually in your arm. A tube collects the blood sample. You may feel a brief scratch or sting.

The blood draw normally takes a few minutes. Afterward, pressure and a small dressing are placed over the site.

The sample is sent to a laboratory. Results are often available within several days, although timing varies by laboratory and whether the sample needs to be sent elsewhere.

Understanding your results

Your result should be compared with the reference range printed on your laboratory report. There is no single range suitable for every person or every testing method. Ranges and interpretation can vary by laboratory, age, sex, pregnancy status, medicines, and clinical context.

A high TBG level means more carrier protein is available. This often raises total T4 and total T3. If free T4 and TSH are normal, it does not necessarily mean your thyroid is overactive.

A low TBG level means less carrier protein is available. This may lower total T4 and total T3 even when the amount of free hormone is appropriate and the thyroid is working normally.

Your TBG result is usually interpreted alongside:

An inherited TBG difference generally affects hormone transport rather than thyroid hormone production. It usually does not require thyroid treatment when TSH and free hormone results show normal thyroid function.

- TSH
- Free T4

- Sometimes total T4, total T3, or free T3
- Your medicines, pregnancy status, and medical history
- Previous results and the laboratory's reference range

What affects the results

Pregnancy commonly raises TBG because estrogen increases the amount of this protein in the blood. Combined contraceptive pills and oral estrogen therapy can have a similar effect. TBG may remain altered for a period after estrogen levels or treatment change.

Androgens, including testosterone and anabolic steroids, can lower TBG. Glucocorticoid medicines may also reduce it. Some medicines change how thyroid hormones attach to carrier proteins without changing the measured TBG concentration, which can make related thyroid results harder to interpret.

Liver conditions can raise or lower TBG because the protein is produced and processed by the liver. Severe illness may also change blood protein levels.

Conditions that cause substantial protein loss in urine, such as nephrotic syndrome, can lower TBG. Reduced kidney function by itself does not always have the same effect; the amount of protein being lost and the wider clinical situation matter.

Inherited changes in the SERPINA7 gene can cause partial or complete TBG deficiency or, less commonly, TBG excess. Some variants change how well TBG binds thyroid hormone rather than simply changing its amount.

Age matters, particularly for newborns and children. Sex-related hormone exposure and pregnancy also affect interpretation. A large recent transfusion or major intravenous fluid replacement may temporarily alter blood protein concentrations, but an ordinary past transfusion is not usually a major influence.

Food, coffee, routine daily activity, and the time of day have little direct effect on TBG. Hard exercise is not considered a major source of misleading TBG results, although acute illness or extreme physical stress can affect the wider thyroid test pattern.

Laboratory interference is uncommon but possible. Antibodies in your blood can sometimes react with an immunoassay and produce a result that does not fit the true TBG level. High-dose biotin can interfere with some laboratory platforms, especially when other thyroid tests are measured at the same time.

Accuracy and limitations

TBG assays generally measure the amount of protein reliably, but the test does not answer whether your thyroid gland is underactive or overactive. A genuinely high or low TBG result can be misleading if it is viewed without TSH and free T4.

There is no useful single figure for how often TBG testing "misses" a thyroid disorder. It is not designed to diagnose one. A normal TBG result does not rule out thyroid disease, and an abnormal result does not prove thyroid disease.

Rare inherited TBG variants may not be fully assessed by a routine quantity measurement. A person can have an unusual protein that is present but does not bind hormone normally. Some laboratory methods may also detect particular variants differently, creating an unexpectedly normal, high, or low result.

If the main question is how well your thyroid works, TSH and free T4 are usually better tests. When ordinary free T4 methods remain inconsistent with the clinical picture, a specialist laboratory may measure free T4 using equilibrium dialysis or ultrafiltration. These methods physically separate free hormone from protein-bound hormone. Genetic testing may occasionally help confirm an inherited TBG difference, but it is not routinely required.

Risks and complications

The risks are those of an ordinary blood draw. You may have brief discomfort, a small bruise, or feel light-headed.

Persistent bleeding, increasing redness, warmth, or swelling at the needle site is uncommon and should receive prompt medical advice.

What happens next

A high or low TBG result usually leads to routine follow-up rather than urgent treatment. Your healthcare professional may review your medicines, pregnancy status, liver and kidney health, and other thyroid test results.

If TSH and free T4 are appropriate, you may not need treatment. The result may simply be recorded so that future total T4 or T3 measurements are interpreted correctly. Repeat testing or a different thyroid hormone test may be used if the results do not fit together.

A TBG result alone is not an emergency finding. Seek emergency care for severe symptoms such as difficulty breathing, chest pain, or fainting that does not quickly resolve, regardless of

the test result. These symptoms require assessment for their own cause.

Common questions

Can I drink coffee before a TBG blood test?

Usually, yes. Coffee does not have a meaningful direct effect on TBG. Avoid it only if another test being taken at the same appointment requires fasting.

Does the time of day matter?

TBG does not usually require collection at a particular time. Timing may matter for another test or medicine being checked alongside it, so follow the instructions for the whole blood test appointment.

Can I exercise beforehand?

Normal activity is unlikely to affect the result. Extreme exercise or acute physical stress can influence other thyroid-related measurements, so mention it if your results are unexpected.

Should I stop my medicines first?

No, not unless the person who prescribed or ordered them tells you to. Estrogen, androgens, glucocorticoids, and some other medicines can affect interpretation, but stopping them without guidance may be unsafe.

Will pregnancy change my result?

Yes. Pregnancy usually raises TBG because estrogen levels increase. Laboratories and healthcare professionals interpret thyroid tests differently during pregnancy, using the full pattern rather than TBG alone.

Does a high or low TBG result mean I need thyroid medicine?

Not by itself. Treatment decisions are based mainly on thyroid function, including TSH and free T4, as well as your symptoms and clinical context.

Sources

- MedlinePlus Medical Encyclopedia: TBG blood test , U.S. National Library of Medicine
- Mayo Clinic Laboratories: Thyroxine-Binding Globulin, Serum , laboratory test overview and interpretation

- American Thyroid Association: Thyroid Function Tests , patient information on total and free thyroid hormones
- Endotext: Defects of Thyroid Hormone Transport in Serum , NCBI Bookshelf
- Pappa T, Ferrara AM, Refetoff S. Inherited defects of thyroxine-binding proteins , Best Practice & Research Clinical Endocrinology & Metabolism , 2015;29(5):735-747